

Instability of speckle patterns in random media with noninstantaneous Kerr nonlinearity

S.E. Skipetrov

*Laboratoire de Physique et Modélisation des Milieux Condensés
CNRS, 38042 Grenoble, France*

Onset of the instability of a multiple-scattering speckle pattern in a random medium with Kerr nonlinearity is significantly affected by the noninstantaneous character of the nonlinear medium response. The fundamental time scale of the spontaneous speckle dynamics beyond the instability threshold is set by the largest of times T_D and τ_{NL} , where T_D is the time required for the multiple-scattered waves to propagate through the random sample and τ_{NL} is the relaxation time of the nonlinearity. Inertial nature of the nonlinearity should complicate the experimental observation of the instability phenomenon.

Accepted for publication in *Optics Letters* © 2003 Optical Society of America, Inc.

Instabilities, self-oscillations and chaos are widely encountered in nonlinear optical systems^{1,2,3}. In particular, the so-called modulation instability (MI), consisting in the growth of small perturbations of the wave amplitude and phase, resulting from the combined effect of nonlinearity and diffraction, can be observed for both coherent^{2,4} and incoherent^{5,6,7,8,9} light. In the latter case, propagation of a powerful, spatially incoherent quasimonochromatic light beam through an optically *homogeneous* nonlinear crystal results in ordered or disordered spatial patterns^{5,6,7,8,9}. Recently, instabilities have also been predicted^{10,11} for diffuse light in *random* media with Kerr nonlinearity (sample size $L \gg$ mean free path $\ell \gg$ wavelength λ). The strength of the nonlinearity has to exceed a threshold for the instability to develop, similarly to the case of incoherent MI (coherent MI has no threshold), but in contrast to the latter, (a) initially coherent wave loses its spatial coherence *inside* the medium, due to the multiple scattering on heterogeneities of the refractive index, (b) temporal coherence of the incident light is assumed to be perfect (in the case of incoherent MI, the coherence time of the incident beam is much shorter than the response time of the medium), (c) incident light beam is completely destroyed by the scattering after a distance $\sim \ell$ and light propagation is diffusive in the bulk of the sample, and (d) scattering provides a distributed feedback mechanism, absent in the case of MI.

Since the noninstantaneous nature of nonlinearity proved to be important for the development of optical instabilities^{1,2,3,4,5,6,7,8,9}, it is natural to consider its effect on the instability of diffuse waves in random nonlinear media. Up to now, the latter phenomenon has only been studied assuming the instantaneous medium response^{10,11}. In the present letter, we show that while the absolute instability threshold does not depend on the nonlinearity response time τ_{NL} , the dynamics of the speckle pattern above the threshold is extremely sensitive to the relation between τ_{NL} and the time $T_D = L^2/D$ that takes the diffuse wave to propagate through the ran-

dom sample ($D = c\ell/3$ is the diffusion constant and c is the speed of light in the average medium). We derive explicit expressions for the characteristic time scale τ of spontaneous intensity fluctuations beyond the instability threshold and the maximal Lyapunov exponent $\Lambda_{\max}$, characterizing the predictability of the system behavior. Finally, we discuss the experimental implications of our results.

We consider a plane monochromatic wave of frequency ω_0 incident on a random sample of typical size $L \gg \ell$. The real dielectric constant of the sample can be written as $\varepsilon(\mathbf{r}, t) = \varepsilon_0 + \delta\varepsilon(\mathbf{r}) + \Delta\varepsilon_{NL}(\mathbf{r}, t)$, where ε_0 and $\delta\varepsilon(\mathbf{r})$ are the average value and the fluctuating part of the linear dielectric constant, respectively, and $\Delta\varepsilon_{NL}(\mathbf{r}, t)$ denotes the nonlinear part of ε . Without loss of generality, we assume $\varepsilon_0 = 1$ and $\langle \delta\varepsilon(\mathbf{r})\delta\varepsilon(\mathbf{r}_1) \rangle = 4\pi/(k_0^4\ell)\delta(\mathbf{r}-\mathbf{r}_1)$. Under conditions of weak scattering $k_0\ell \gg 1$ (where $k_0 = \omega_0/c$), the average intensity $\langle I(\mathbf{r}) \rangle$ of the scattered wave inside the sample can be described by a diffusion equation, the short-range correlation function of intensity fluctuations $\delta I(\mathbf{r}, t) = I(\mathbf{r}, t) - \langle I(\mathbf{r}) \rangle$ decreases to zero at distances of order λ , and the long-range correlation of $\delta I(\mathbf{r}, t)$ can be found from the Langevin equation¹² (see Ref. 13 for a review of wave diffusion in random media):

$$\frac{\partial}{\partial t}\delta I(\mathbf{r}, t) - D\nabla^2\delta I(\mathbf{r}, t) = -\nabla \cdot \mathbf{j}_{\text{ext}}(\mathbf{r}, t). \quad (1)$$

In a nonlinear medium, the random external Langevin currents $\mathbf{j}_{\text{ext}}(\mathbf{r}, t)$ obey a dynamic equation¹¹

$$\begin{aligned} \frac{\partial}{\partial t}\mathbf{j}_{\text{ext}}(\mathbf{r}, t) &= \int_V d^3\mathbf{r}' \int_0^\infty d\Delta t \mathbf{q}(\mathbf{r}, \mathbf{r}', \Delta t) \\ &\times \frac{\partial}{\partial t}\Delta\varepsilon_{NL}(\mathbf{r}', t - \Delta t), \end{aligned} \quad (2)$$

where the integration is over the volume V of the random medium and $\mathbf{q}(\mathbf{r}, \mathbf{r}', \Delta t)$ is a random, sample-specific response function with zero average and a correlation function given by the diagrams of Fig. 1.

Supplemented by a Debye-relaxation relation for




FIG. 1: Diagrams contributing to the correlation function $\langle q^{(i)}(\mathbf{r}, \mathbf{r}', t - t') q^{(j)*}(\mathbf{r}_1, \mathbf{r}'_1, t_1 - t'_1) \rangle$ of the random response functions $\mathbf{q}$ entering into Eq. (2). Solid lines represent the retarded and advanced Green's functions of the linear wave equation, dashed lines denote scattering of the two connected wave fields on the same heterogeneity. Wavy lines are k_0^2 vertices.

$\Delta \varepsilon_{\text{NL}}(\mathbf{r}, t)$:

$$\tau_{\text{NL}} \frac{\partial}{\partial t} \Delta \varepsilon_{\text{NL}}(\mathbf{r}, t) = -\Delta \varepsilon_{\text{NL}}(\mathbf{r}, t) + 2n_2 I(\mathbf{r}, t), \quad (3)$$

where n_2 is the nonlinear coefficient, Eqs. (1) and (2) can now be used to perform a linear stability analysis of the multiple-scattering speckle pattern in a random medium with noninstantaneous nonlinearity. Assuming $\delta I(\mathbf{r}, t) = \delta I(\mathbf{r}, \alpha) \exp(\alpha t)$ with $\alpha = i\Omega + \Lambda$ ($\Omega > 0$), and similarly for $\mathbf{j}_{\text{ext}}(\mathbf{r}, t)$ and $\Delta \varepsilon_{\text{NL}}(\mathbf{r}, t)$, we use Eqs. (2) and (3) to express the correlation function of Langevin currents $\langle \mathbf{j}_{\text{ext}}^{(i)}(\mathbf{r}, \alpha) \mathbf{j}_{\text{ext}}^{(j)*}(\mathbf{r}_1, \alpha) \rangle$ in terms of the intensity correlation function $\langle \delta I(\mathbf{r}, \alpha) \delta I^*(\mathbf{r}_1, \alpha) \rangle$ and Eq. (1) to express the latter correlation function in terms of the former. For large enough sample size $L/\ell \gg k_0 \ell$ and moderate frequency $\Omega T_D \ll [L/(k_0 \ell^2)]^2$ we can neglect the short-range correlation of intensity fluctuations and write the condition of consistency of the two obtained equations [and hence the condition of consistency of Eqs. (1-3)] as

$$p \simeq F(\Omega T_D, \Lambda T_D) H(\Omega \tau_{\text{NL}}, \Lambda \tau_{\text{NL}}), \quad (4)$$

where a numerical factor of order unity has been omitted, $p = \langle \Delta n_{\text{NL}} \rangle^2 (L/\ell)^3$ is the effective nonlinearity parameter, $\langle \Delta n_{\text{NL}} \rangle = n_2 \langle I(\mathbf{r}) \rangle$ is the average value of the nonlinear part of refractive index, and $\langle I(\mathbf{r}) \rangle$ is assumed to be approximately constant inside the sample. The function F in Eq. (4) is the same as in the case of instantaneous nonlinearity¹¹, while $H(x, y) = x^2 + (1 + y)^2$ originates from the noninstantaneous nature of the nonlinear response. In the limit of $\tau_{\text{NL}} \rightarrow 0$ (instantaneous nonlinearity), $H \equiv 1$ and Eq. (4) reduces to Eq. (7) of Ref. 11.




FIG. 2: Instability threshold as a function of the excitation frequency Ω (in units of T_D^{-1}) for several ratios of the nonlinearity relaxation time τ_{NL} and the time T_D required for light to diffuse through the random sample. Dashed line corresponds to $\tau_{\text{NL}}/T_D = 0$. Inset: Lyapunov exponent Λ vs. Ω (both in units of T_D^{-1}) slightly above the threshold ($p = 1.1$) for the same values of τ_{NL}/T_D as in the main plot.

For given Ω and p , Eq. (4) determines the corresponding value of the Lyapunov exponent Λ . $\Lambda > 0$ signifies the instability of the speckle pattern with respect to the excitations at frequency Ω . If $p < 1$, then $\Lambda < 0$ for all Ω . As p exceeds 1, Λ becomes positive inside some frequency band $(0, \Omega_{\text{max}})$ (see the inset of Fig. 2). We show the frequency-dependent threshold following from Eq. (4) in Fig. 2 for different values of τ_{NL}/T_D . The threshold value of p at $\Omega \gtrsim \tau_{\text{NL}}^{-1}$ deviates from the result obtained for the instantaneous nonlinearity (dashed line in Fig. 2), while the *absolute* threshold is $p = 1$ at $\Omega = 0$, independent of τ_{NL}/T_D .

To study the onset of the instability, we consider the limit of $0 < p - 1 \ll 1$ in more detail. In this limit, a series expansion of the function F in Eq. (4) can be found: $F(x, y) \simeq 1 + C_1 x^2 + y(C_2 - C_3 x^2)$, where we keep only the terms quadratic in x and linear in y , and the numerical constants are $C_1 \simeq 0.031$, $C_2 \simeq 0.59$, $C_3 \simeq 0.045$. At a given frequency Ω , the threshold value of p then becomes $p \simeq 1 + [C_1 + (\tau_{\text{NL}}/T_D)^2] (\Omega T_D)^2$, while at a given p the maximum excited frequency is $\Omega_{\text{max}} = T_D^{-1} [C_1 + (\tau_{\text{NL}}/T_D)^2]^{-1/2} (p - 1)^{1/2}$ and the shortest typical time of spontaneous intensity fluctuations is $\tau = 1/\Omega_{\text{max}}$.

As follows from the above analysis, a continuous low-frequency spectrum of frequencies $(0, \Omega_{\text{max}})$ is excited at $p > 1$, and the Lyapunov exponent Λ decreases monotonically with Ω (see the inset of Fig. 2). This allows us to hypothesize that at $p = 1$ the speckle pattern undergoes a transition from a stationary to chaotic state. Such a behavior should be contrasted from the “route to chaos” through a sequence of bifurcations, characteristic of many nonlinear systems^{1,2,3}. Given initial conditions at $t = 0$ and deterministic dynamic equations, the

evolution of a chaotic system can only be predicted for $t < 1/\Lambda_{\max}$. We find the maximal Lyapunov exponent $\Lambda_{\max} = T_D^{-1}(p-1)/(C_2 + 2\tau_{\text{NL}}/T_D)$. It is now easy to see that the fundamental time scale of the speckle pattern dynamics at $p > 1$ is set by the largest of the times T_D and τ_{NL} . Indeed, in the case of fast nonlinearity ($\tau_{\text{NL}} \ll T_D$) we have $\Omega_{\max} \propto T_D^{-1}(p-1)^{1/2}$, $\tau \propto T_D(p-1)^{-1/2}$, and $\Lambda_{\max} \propto T_D^{-1}(p-1)$. In the opposite limit of slow nonlinear response ($\tau_{\text{NL}} \gg T_D$) we find $\Omega_{\max} \propto \tau_{\text{NL}}^{-1}(p-1)^{1/2}$, $\tau \propto \tau_{\text{NL}}(p-1)^{-1/2}$, and $\Lambda_{\max} \propto \tau_{\text{NL}}^{-1}(p-1)$, respectively.

The origin of the instability of the speckle pattern $I(\mathbf{r}, t)$ in a nonlinear random medium and its sensitivity to τ_{NL} can be qualitatively understood by considering I as a result of interference of many partial waves traveling along various diffusion paths. An infinitely small perturbation of I at some point $\mathbf{r}_0$ at time t_0 diffuses throughout the medium [according to Eq. (1)] and, after some time Δt , modifies the speckle pattern inside a sphere of radius $R \sim (D\Delta t)^{1/2}$ around $\mathbf{r}_0$. Due to the nonlinearity of the medium, this leads to a change of the nonlinear part of the dielectric constant $\Delta\epsilon_{\text{NL}}$ inside the same sphere after a time $\sim \tau_{\text{NL}}$ [according to Eq. (3)] and, consequently, to a modification of the phases of the partial diffuse waves that interfere to produce a new speckle pattern $I(\mathbf{r}, t)$. If the strength of the nonlinearity exceeds the threshold given by Eq. (4), $I(\mathbf{r}_0, t) - I(\mathbf{r}_0, t_0)$ can be larger than the initial perturbation. The latter is hence amplified and the speckle pattern develops spontaneous dynamics. Noninstantaneous nature of the nonlinearity leads to the damping of the high-frequency components in the spectrum of the initial intensity perturbation, raising the

instability threshold for $\Omega \gtrsim \tau_{\text{NL}}^{-1}$, as we show in Fig. 2. This phenomenon is similar in its origin to the increase of the speed of coherent MI patterns with decreasing τ_{NL} in a *homogeneous* nonlinear medium⁴. Similarly to the case of coherent MI⁴, the instability is arrested in the limit of $\tau_{\text{NL}} \rightarrow \infty$, in contrast to the case of incoherent MI, where $\tau_{\text{NL}} \rightarrow \infty$ (more precisely, $\tau_{\text{NL}} \gg$ than the coherence time of the incident beam) is an explicit assumption of the theoretical models^{5,6,9}.

To conclude, we discuss the experimental implications of our results. Δn_{NL} up to 10^{-3} can be realized in inorganic photorefractive crystals^{6,7,8}. If scattering centers (optical defects) are introduced in the crystal, $L/\ell \sim 10^2$ will suffice to reach the instability threshold $p \simeq 1$. Alternatively, in nematic liquid crystals both nonlinearity (Δn_{NL} up to 0.1, see, e.g., Ref. 14) and scattering ($L/\ell > 10$ has been reported in Ref. 15) can be sufficiently strong to obtain $p \simeq 1$. The nonlinearity is rather slow in the above cases, τ_{NL} exceeds T_D ($T_D \sim 10 \div 100$ ns for $\ell \sim 0.1 \div 1$ mm and $L \sim 1$ cm), and the results obtained in the present letter are, therefore, of particular relevance. Finally, although the noninstantaneous nature of nonlinearity does not affect the absolute instability threshold, it should complicate the experimental observation of the instability phenomenon. Indeed, in a real experiment only sufficiently fast fluctuations can be detected. Inasmuch as $\Omega_{\max}$ decreases with τ_{NL} , larger p (and thus stronger nonlinearities) will be required for the instability to be detectable in a medium with noninstantaneous nonlinearity.

-
- ¹ H.M. Gibbs, “Optical Bistability: Controlling Light With Light” (Academic Press, Orlando, 1985).
 - ² G.P. Agrawal, “Nonlinear Fiber Optics” (Academic Press, San Diego, 1995).
 - ³ F.T. Arecchi, S. Boccaletti, P. Ramazza, “Pattern formation and competition in nonlinear optics”, Phys. Rep. **318**, 1–83 (1999).
 - ⁴ M.-F. Shih, C.-C. Jeng, F.-W. Sheu, and C.-Y. Lin, “Spatiotemporal optical modulation instability of coherent light in noninstantaneous nonlinear media”, Phys. Rev. Lett. **88**, 133902 (2002).
 - ⁵ M. Soljacic, M. Segev, T. Coskun, D.N. Christodoulides, and A. Vishwanath, “Modulation instability of incoherent beams in noninstantaneous nonlinear media”, Phys. Rev. Lett. **84**, 467–470 (2000).
 - ⁶ D. Kip, M. Soljacic, M. Segev, E. Eugenieva, and D.N. Christodoulides, “Modulation instability and pattern formation in spatially incoherent light beams”, Science **290**, 495–498 (2000).
 - ⁷ J. Klinger, H. Martin, and Z. Chen, “Experiments on induced modulational instability of an incoherent optical beam”, Opt. Lett. **26**, 271–273 (2001).
 - ⁸ Z. Chen, S.M. Sears, H. Martin, D.N. Christodoulides, and M. Segev, “Clustering of solitons in weakly correlated wavefronts”, Proc. Nat. Acad. Sci. **99**, 5223–5227 (2002).

- ⁹ B. Hall, M. Lisak, D. Anderson, R. Fedele, and V.E. Semenov, “Statistical theory for incoherent light propagation in nonlinear media”, Phys. Rev. E **65**, 035602 (2002).
- ¹⁰ S.E. Skipetrov and R. Maynard, “Instabilities of waves in nonlinear disordered media”, Phys. Rev. Lett. **85**, 736–739 (2000) (see also cond-mat/0006136).
- ¹¹ S.E. Skipetrov, “Langevin description of speckle dynamics in nonlinear disordered media”, Phys. Rev. E **67**, 016601 (2003) (see also cond-mat/0206475).
- ¹² A.Yu. Zyuzin and B.Z. Spivak, “Langevin description of mesoscopic fluctuations in disordered media” Sov. Phys. JETP **66**, 560–566 (1987).
- ¹³ M.C.W. van Rossum and Th.M. Nieuwenhuizen, “Multiple scattering of classical waves: microscopy, mesoscopy, and diffusion”, Rev. Mod. Phys. **71**, 313–371 (1999).
- ¹⁴ R. Muenster, M. Jarasch, X. Zhuang, and Y.R. Shen, “Dye-induced enhancement of optical nonlinearity in liquids and liquid crystals”, Phys. Rev. Lett. **78**, 42–45 (1997).
- ¹⁵ M.H. Kao, K.A. Jester, A.G. Yodh, and P.J. Collings, “Observation of light diffusion and correlation transport in nematic liquid crystals”, Phys. Rev. Lett. **77**, 2233–2236 (1996).